

## REVIEW

# *Can Amiodarone Prevent Sudden Cardiac Death in Patients with Hemodynamically Tolerated Sustained Ventricular Tachycardia and Coronary Artery Disease?*

Arash Arya, Majid Haghjoo, and Mohammad Ali Sadr-Ameli

Department of Pacemaker and Electrophysiology, Rajaie Cardiovascular Medical Center, Mellat Park, Vali-Asr Avenue, Tehran, 1996911151, Iran

**Summary.** One of the most important challenges in today's cardiology is prevention of sudden cardiac death in high risk patients with coronary artery disease (CAD). Sustained hemodynamically tolerated ventricular tachycardia (HTVT) comprises up to 30% of all cases of monomorphic ventricular tachycardia in patients with CAD. While there is a consensus on treatment of hemodynamically unstable sustained ventricular tachycardia in patients with CAD, some controversies regarding the proper treatment of HTVT exist. We re-examined existing clinical evidence, controversies and current guidelines on the treatment of HTVT in patients with CAD and demonstrated that compared to implantable cardioverter-defibrillator, amiodarone is not an acceptable therapeutic option in patients with ischemic heart disease who suffer from HTVT.

**Key Words.** coronary artery disease, ventricular tachycardia, implantable defibrillators, amiodarone, sudden cardiac death, secondary prevention

## Introduction

One of the most important challenges nowadays is prevention of sudden cardiac death (SCD) in high risk patients with coronary artery disease (CAD) [1]. There is some inconsistent evidence supporting potential benefit of amiodarone in the prevention of SCD in high risk post myocardial infarction (MI) patients [2]. Development of implantable cardioverter-defibrillators (ICD) has been a dramatic advancement in the management of patients with ventricular tachycardia (VT). Several reviews have assessed the current evidence on superiority of ICD in the prevention of SCD in various patient populations with spontaneous sustained monomorphic ventricular tachycardia (MMVT) compared to amiodarone, and, based on these studies, AHA/ACC guideline has given ICD a class I indication with level of evidence: B in patients with spontaneous sustained VT (irrespective of hemodynamic status during arrhythmia) in association with structural heart disease [3,4]. While there is a consensus on treatment of hemodynamically unstable sustained VT in patients with CAD, some controversies exist regarding the proper treatment of HTVT [1,4]. This review intends to re-examine existing

clinical evidence, controversies and current guidelines on the treatment of HTVT in patients with CAD.

## *Prevalence of MMVT and HTVT in patients with CAD and its impact on survival*

Late sustained MMVT occurs in 3–5% [5] of patients after an acute MI and has been associated with a poor prognosis (relative risk of mortality: 2.6 to 9.1 according to different studies) when compared to those without a ventricular arrhythmia [6]. Several studies have assessed the effect of MMVT on survival of patients with CAD [6–10,13]. The reported annual mortality of these patients varied from 5% (in those with a left ventricular ejection fraction [LVEF] >50%) [9] to more than 40% [10] in patients with LV dysfunction. Newby et al. [7] have examined the incidence and impact of MMVT on survival of 40,895 post MI patients in the “Global Use of Streptokinase t-PA for Occluded Coronary Arteries” (GUSTO)-I trial. In GUSTO-I the incidence of late sustained MMVT was 3.5%. The overall one year mortality in 30 day survivors in the late VT group was 24.7% (mean LVEF = 46%) compared to 2.7% in patients without ventricular arrhythmias (mean LVEF = 52%). Al-Khatib et al. [11] have recently assessed the effect of late MMVT on survival of 15,042 post MI patients participating in GUSTO-III trial which confirmed the above mentioned findings of GUSTO-I trial. These results were confirmed also by a study on 26,416 patients with acute coronary syndrome [12]. The prevalence of late MMVT in these patients was 2.1% (lower than post MI patients in GUSTO-I and III), however the hazard ratio (HR) was comparable (HR = 5, 95% Confidence Interval = 3.8–6.5) to the GUSTO-I and III studies. It is worth to mention that in GUSTO I and III the mortality of patients with late

Address for correspondence: Arash Arya, M.D., Department of Pacemaker and Electrophysiology, Rajaie Cardiovascular Medical Center, Mellat Park, Vali-Asr Avenue, Tehran 1996911151, Iran. Tel.: 0098 912 126 1467; Fax: 0098 21 878 4618; E-mail: arya@rhc.ac.ir

sustained MMVT was higher compared to patients with late VF. This is in accordance to the CARE group results [14] which showed that the probability of appropriate ICD discharge is two times higher in patients with sustained MMVT compared to those with aborted SCD and VF.

A recently published guideline [1] has recommended amiodarone and beta blocker therapy as a class IIa indication and ICD, along with ablation and surgery, as class IIb recommendation for the treatment of patients with HTVT. The above mentioned statement could only be accepted if one assumes that the mortality in HTVT is significantly lower than more severely symptomatic VTs and that amiodarone therapy is equal or superior to ICD with respect to prevention of SCD in this particular group of patients.

HTVT comprises up to 30% of all cases of MMVT in patients with CAD [15,16]. Several studies have examined the effect of HTVT on survival compared to more severely symptomatic VT. Although Sarter et al. have suggested a better prognosis for HTVT [17] (however, some debates exist on their data as 64% of deaths in their study were non-sudden either due to perioperative death, recurrent infarction or progressive heart failure. Thirty seven percent of patients were treated with VT surgery with a perioperative mortality of 20%. This surgery improved the outcome in patients who survived the operation and gives an inaccurate estimate of the risk of SCD in those who survived. In addition they found that longer VT cycle length, which one could expect to be associated with more benign symptoms, is related to a higher mortality [18]). Others showed that the risk is similar to patients with more severely symptomatic VT. Raitt et al. performed a retrospective subgroup analysis of the AVID registry [15] and showed that the absence of symptoms with sustained VT does not predict a benign prognosis (see below). Olson et al. assessed the predictors of SCD in 122 patient followed for an average of 19.5 months [18] and showed that the rate of SCD is not affected by presence or absence of symptoms during MMVT.

Multiple VTs (including very rapid, poorly tolerated VTs) are commonly induced during electrophysiological (EP) testing in patients with stable VT [19]. Having these in mind, HTVT actually is a marker of substrates for re-entrant ventricular arrhythmias which may causes more malignant ventricular arrhythmias during *long term* follow up. Based on available data, ICD therapy decreases all-cause mortality in CAD patients with sustained VT, and with respect to the above mentioned findings, could also decrease the mortality in patients with HTVT. Patients with HTVT are at high risk for sudden arrhythmic death, and presumably it is not the recurrence of the stable VT that leads to SCD but a more malignant ventricular arrhythmias. Bocker et al. studied the natural course of 50 patients (82%, CAD) with HTVT who received an ICD [20]. They showed that during mean follow up of 17 months, 33 patients (66%) had 3861 episodes of ventricular tachycar-

dia which is comparable to other studies in patients with sustained MMVT [14,21]. Ninety one percent of these episodes were terminated by antitachycardia pacing. Eleven patients (22%) had episodes of potentially life-threatening fast VT (CL <250 ms) during follow up period. In the AVID registry [18] (mean follow up of 16.9 months) the mortality rate for patients with syncope VT was 21.2% and for asymptomatic VT was 19.7% ( $P = \text{NS}$ ). Had the ICD not been implanted in the Bocker study, their patients would have had at least the same mortality as in the AVID registry. It is worth to mention that in the Bocker study (like the Electrophysiologic Study Versus Electromagnetic Monitoring trial) [22] EP studies failed to predict which patient would have more rapid VT in the follow up. In conclusion, currently available data depict that HTVT negatively affects the survival of post MI patients as more severely symptomatic VT does.

Finally, epidemiological studies have consistently shown an inverse relationship between LVEF and survival in patients with CAD [23]. When LVEF decreases below 35–40%, the risk of sudden and non-sudden cardiac death rises sharply. In patients who have an LVEF <40%, the mortality rate at 2 years after infarction is 21.5% compared with 7.6% in patients with an LVEF >40% [23–27].

A subgroup analysis of AVID showed that patients with LVEF >35% failed to benefit from ICD [28]. However, the AVID trial design had a low power to detect an ICD benefit in this small subgroup of patients during short observation period which was also terminated prematurely. These factors prevented the ICD benefit from reaching statistical significance in this small subgroup of patients. As we indicate below (see: Do we need a prospective clinical trial comparing ICDs with empiric or EP guided amiodarone therapy in CAD patients with HTVT) if we want to conduct an AVID like trial in patients with HTVT and preserved LV function one would need a sample size of 1072 patients followed for three years. Hence, had the follow up period of the AVID trial been continued in an adequate number of patients for a sufficient time period, ICD would have shown its beneficial effect in these patients (see also: Do the benefits of ICD versus Amiodarone change over time?) In addition there is another subgroup analysis of AVID which showed that in patients treated with antiarrhythmic medications (but not those treated with ICD) survival was strongly associated with LVEF. The lack of statistically significant association in ICD patients might have been related to its efficacy in terminating malignant VA *regardless* of LVEF [29].

## ***Role of Amiodarone in the Treatment of Patients with Sustained VT***

### ***A: Empiric Amiodarone Therapy***

*Secondary prevention.* Amiodarone suppresses premature ventricular depolarizations and episodes of

nonsustained VT, but, there have not been any placebo-controlled trials on its effectiveness on sustained VT and VF [2]. All available articles only report the outcomes of patients with aborted sudden cardiac arrest or recurrent VT treated with amiodarone alone or versus other antiarrhythmics. Some reports conclude that amiodarone is as effective, and some suggest that amiodarone is not as effective as it was shown by early promising reports. The Cardiac Arrest in Seattle: Conventional versus Amiodarone Drug Evaluation study, the only randomized clinical trial available, showed that amiodarone is superior to other conventional antiarrhythmic drugs, which we know today, increase cardiac mortality [30]. The largest follow-up of amiodarone treated patients [31] (589 patients with supraventricular tachycardia, 83% of whom had VT or VF), showed that the 5-year cumulative risk of sudden death was 22% and of total mortality, 46%. The cumulative risk of drug failure (defined as SCD, recurrence of ventricular arrhythmias, or drug withdrawal) at 5 years, was 50%. In conclusion it is difficult to reach to any definite conclusion about the efficacy of amiodarone from these uncontrolled reports.

**Primary prevention.** Fifteen randomized clinical trials (5864 patients), including six in post MI patients [32–37], were performed on amiodarone as a prophylaxis against SCD in moderate to high risk patients for SCD. Two meta-analyses [37,38] of these trials showed a 13 to 19% reduction in total mortality but the odds ratio was different based on the control group: the odds ratio for total mortality was lower in trials with “usual care” controls (odds ratio, 0.58; 95% CI, 0.41 to 0.83;  $P = .003$ ) and in trials with active controls (odds ratio, 0.73; 95% CI, 0.43 to 1.25;  $P = .25$ ) than in trials with placebo controls (odds ratio, 0.90; 95% CI, 0.76 to 1.06;  $P = .20$ ). These two meta-analyses suggested that amiodarone therapy reduces total mortality by between 10% (placebo-controlled trials only,  $P = \text{NS}$ ) and 13–19% (all trials,  $P = 0.03$  and  $P < 0.01$  respectively) in patients with moderate to high risk of sudden cardiac death. There has been no placebo-controlled trial so far to assess amiodarone’s effect in patients with HTVT.

Finally, the results of The Sudden Cardiac Death in Heart Failure Trial (SCD-HeFT) confirmed the findings in the above mentioned placebo-controlled studies on amiodarone. Among patients with NYHA class II and III congestive heart failure and an LVEF  $\leq 35\%$  (23% with a history of NSVT) who were on optimal medical therapy, amiodarone (compared to placebo) did not show a beneficial effect on total mortality by intention to treat analysis (HR = 1.06, 97.5% Confidence Interval = 0.86–1.30,  $P = 0.529$ ) [39].

### **B: EP-guided amiodarone therapy**

Amiodarone is usually prescribed empirically (as it is recommended by ESC taskforce on SCD). Several studies have suggested that EP-guided therapy can in-

crease the success rate of therapy with amiodarone [40–47]. In these studies lack of inducible ventricular arrhythmias after amiodarone therapy was associated with a better outcome. Lack of suppression of ventricular arrhythmias, slowing of the VT cycle length, and unchanged ventricular effective refractory period were all associated with higher long term recurrence of ventricular arrhythmias and mortality. There are three major setbacks in EP-guided amiodarone therapy. First, the success rate for complete VT suppression rate during EP studies varies between 10–40% in these studies. Second, there is no standard and widely accepted ventricular stimulation protocol for assessment of its usefulness and different protocols have been used so far. Third, amiodarone has not been tested against ICD in the above mentioned trials.

Schl pfer et al. conducted the first study aiming at a comparison between EP-guided amiodarone and ICD therapy in 84 consecutive post MI patients with sustained MMVT [16]. Aborted SCD and syncope were clinical presentations of the index arrhythmia in 40% of cases and 77% of their patients were in NYHA class = II. They showed that the outcome of the patients (including 55% with EF  $\leq 35\%$ ) in their study was better with ICD than EP-guided amiodarone therapy. During follow up of  $63 \pm 30$  months, total mortality (and SCD) was 42% (21%) in EP-guided group and 15% (2%) in ICD group. It is noteworthy that their data showed that even *complete* suppression of VT by EP-guided amiodarone therapy was not protective against risk of future SCD (Schl pfer J: Personal communication).

### **C: adjunctive amiodarone in patients with ICD**

No empiric antiarrhythmic therapy (including amiodarone) is currently indicated in patients who received an ICD. Up to 40% of patients receiving an ICD develop “electrical storm,” defined as two or more episodes of VT and/or VF in a one day period [48,49]. These patients frequently receive multiple ICD shocks, which severely impair quality of life. Intravenous followed by oral amiodarone results in successful management and possibly a long-term effect similar to patients who do not have electrical storm [48,49].

The OPTIC (Optimal Pharmacological Therapy in Implantable Cardioverter) study currently assesses the potential benefit of antiarrhythmic medications in reduction of ICD therapy and electrical storm. In OPTIC the patients are randomized to  $\beta$ -blocker, amiodarone plus  $\beta$ -blocker, or sotalol. A sub-study of the OPTIC study will also assess defibrillation threshold before and after drug therapy in patients randomized to the above mentioned drugs [48]. Amiodarone may have some other potential benefits in patients with ICDs including the prevention of supraventricular tachyarrhythmias, which could cause inappropriate ICD discharges; and the prevention of nonsustained but

symptomatic ventricular arrhythmias. Further studies are warranted to clarify this issue [50].

#### **D: role of amiodarone in hybrid therapy of HTVT**

Although several studies have established the role of catheter ablation of VT (including HTVT) as an adjunctive therapy to ICD in patients with CAD who receive frequent high voltage therapies [51–54], few have evaluated the role of catheter ablation plus adjunctive amiodarone in patients with HTVT [55].

Della Bella et al. in a non-randomized prospective study (median follow up of 41.5 months), evaluated the outcome of catheter ablation and adjunctive amiodarone and/or beta-blocker therapy in 124 (mean age  $64 \pm 9$  years, with EF  $>30\%$  in 2/3 of cases) post-MI patients with HTVT (24 of whom had an ICD). The procedure of catheter ablation was successful (defined as termination and prevention of induction of all clinical and inducible VTs) in 73% of patients and after ablation 86% of patients received amiodarone and/or a beta blocker as an adjunctive therapy. The rate of all cause mortality and sudden cardiac death during the study period were 12 and 2.6%, respectively [55]. However, several points merit consideration. Although the mortality was relatively low during follow up, the rate of recurrent VT remained high even among patients with an initially successful procedure. Among these patients the 1 and 3 year VT recurrence rates were 19 and 27%, respectively. In addition, as this study was non-randomized we cannot conclude that this combination therapy is as effective as or superior to ICD in patients with HTVT [55].

In conclusion, based on currently available data [18,20,52] catheter ablation (with or without adjunctive antiarrhythmic therapy) can not be recommended as an alternative to ICD in patients with HTVT. However, in case of frequent ICD shocks, amiodarone and/or catheter ablation should be considered as an adjunctive treatment.

#### **Do the benefits of ICD versus amiodarone change over time?**

A meta-analysis of the CASH, CIDS and AVID trials showed a significant reduction in death from any cause with ICD, hazard ratio (ICD:amiodarone) of 0.72 (95% CI, 0.60 to 0.87;  $P = 0.0006$ ) [56]. However, neither the CIDS nor the CASH trials alone demonstrated a significant benefit of the ICD over amiodarone. Bokhari et al. [57] have recently published the 11 year follow up in a subset of patients of CIDS trial. After a mean follow-up of  $5.6 \pm 2.6$  years in 120 patients, there were 28 deaths (47%) in the amiodarone group, compared with 16 deaths (27%) in the ICD group ( $P = 0.0213$ ). Total mortality was 5.5% per year in the amiodarone group versus 2.8% per year in the ICD group (hazard ratio of amiodarone: ICD, 2.011; 95% confidence interval, 1.087 to 3.721;  $P = 0.0261$ ). In the amiodarone

group, 49 patients (82% of all patients) had side effects related to amiodarone, of which 30 patients (50% of all patients) required discontinuation or dose reduction; 19 patients crossed over to ICD because of amiodarone failure ( $n = 7$ ) or side effects ( $n = 12$ ) [57]. The authors showed that during long term follow up the benefit of the ICD over amiodarone increases and most amiodarone-treated patients eventually develop side effects, have arrhythmia recurrences, or die. This finding has been confirmed recently by Salukhe et al. [58]. They estimated, from published data of 8 major ICD trials, the cumulative benefit life-years gained and calculated the dependency of the benefit on duration of follow-up. They found that the number of life-years gained from one device implantation increases with length of follow-up. Importantly, this increase is markedly *nonlinear*. Within a 3-year span, the benefit rises with the square of time (gain  $\propto t^{1.94}$ ,  $R^2 = 0.998$ ,  $P = 0.001$ ). They concluded that the expected benefit in life span (life-years gained) for a patient who has an ICD is dramatically dependent on the time window over which the benefit is assessed. It is important to consider the effect of follow-up duration while interpreting the results and outcome in ICD trials [58].

Concerns have recently been raised about the role of ICD therapy in apparently stable patients with left ventricular dysfunction several years after MI [59]. It is widely believed that among patients with CAD as the time passes from MI the risk of SCD, and hence, the potential benefit of ICD over amiodarone is diminishing. Long-term follow-up of MI survivors conducted in the 1970s and 1980s indicated that the greatest risk of sudden death was in the initial 6 to 12 months after infarction, particularly in high-risk subgroups such as those with impaired ventricular function [60–62].

Wilber and his colleagues analyzed the time dependence of mortality risk after MI in the MADIT II cohort and evaluated whether long-term survival benefit diminished as a function of elapsed time from infarction to ICD implantation [63]. They found that in contrast to early reports, mortality risk in the MADIT II cohort did not diminish as a function of time from MI; instead, it actually increased. In addition, the survival benefit associated with ICDs appears to be greater for remote MI and remains substantial for up to  $\geq 15$  years after MI. They also found a trend toward increasing device benefit with remote MI although this did not reach statistical significance [63]. In conclusion the above mentioned studies have shown that the benefit of ICD over amiodarone increases over time (Fig. 1). This effect is observed in both primary and secondary prevention trials.

#### **Do we need a prospective clinical trial comparing ICDs with empiric or EP guided amiodarone therapy in CAD patients with HTVT?**

Although there has been no clinical trial (as a gold standard in medical practice) specifically conducted in


patients with HTVT, all the above mentioned data suggest that CAD patients presenting with HTVT have at least as high a total mortality rate as patients presenting with more severely symptomatic VT, and that amiodarone (empiric or EP guided) is not an appropriate option for treatment of patients with sustained MMVT, and an ICD is the treatment of choice similar to patients with unstable VT.

It is sometimes not possible and/or practical to perform a randomized controlled study in every subset of patients at risk [64]. When we have enough reliable data to calculate the risk of adverse event of interest in patients at risk, and a reliable assessment of the beneficial effect of intervention, risk modelling may be used as an alternative [64]. Exner and Klein have suggested that based on available data we can extrapolate the results of available trials with ICD and there is no need to perform a clinical trial in all subsets of patients at high risk for SCD [64]. They estimated that at the completion of the follow up period of SCD-HeFT the ICD would show a statistically significant absolute risk reduction in total mortality (compared to placebo) of 7.3–8.6%, which is confirmed by results of SCD-HeFT [39]. At the completion of the follow up period, ICD resulted in nearly 8% absolute risk reduction in total mortality compared to placebo.

Suppose that we want to assess the effect of ICD vs. amiodarone on total mortality (as primary end point) in patients with HTVT with a clinical trial having a 3 year follow up (Inclusion is 2 years +2 years follow up period after inclusion) and a power of 95% with no interim analysis. We estimate that the total mortality rate would be around 9%/year in the amiodarone group and ICD would reduce the total annual mortality by 50% (the mode of death in patients with HTVT is arrhythmic in at least 61% of cases [63–67] and ICD will effectively terminate these arrhythmias in more than 90% of cases [68], so one can expect that ICD would reduce the total annual mortality by at least 50%) compared to the control group. Based on these assumptions and for a 5% significance level, 2-sided log-rank test of equality of survival curves, 89 events are required during the follow up period which corresponds to a sample size of 536 patients randomized to ICD or amiodarone (Fig. 1). If we want to conduct such a trial only in those patients who have preserved LV function with an estimated annual mortality of  $\approx 5\%$ , the calculated number will be nearly double. We however believe that this trial is not necessary as the bulk of evidence (see above) supports the superiority of ICD in patients who suffer from HTVT, and there would be serious ethical consideration in conducting such a clinical trial.

## Conclusion

Despite current controversies and differences, the available data show that CAD patients with HTVT have a similar prognosis as more severely symptomatic



**Fig. 1.** Survival curves<sup>¶</sup> during hypothetical 5 year follow up of in a low risk group of patients (e.g. those with preserved LV function) treated with ICD vs. amiodarone and control group\*. (\*The annual all cause mortality assumed to be 9% in AMD group and ICD expected to decrease all-cause mortality by 50%. If we randomize 536 patients equally to ICD and AMD group, at the end of three year follow up 89 events would happen and the difference would reach to statistical significance by a 95% power. <sup>†</sup>The calculated survival is based on 10% reduction in all-cause mortality by AMD. <sup>‡</sup>The calculated survival is based on 20% reduction in all-cause mortality by AMD, <sup>¶</sup>Note that survival curves diverge dramatically after second year of follow up and reach to a statistically significant difference after three years of follow up (see above). Had the follow up been stopped after a two years period (like AVID), the difference in mortality would have not reached to statistical significance. This example signifies the effect of follow up duration on assessment of treatment outcome in ICD trials (see also: Does the ICD Benefit Change over Time?)) AMD: Amiodarone.

VT and therefore amiodarone is not an acceptable treatment in the ICD era in these patients. ICD is the preferred mode of therapy in this setting.

## Acknowledgment

The authors would like to thank Elham Hasheminejad, M.D. and Babak Kazemi, M.D. for reviewing of the manuscript. The authors would also like to thank kristof DAEMS for the sample size calculations used in this article.

## References

1. Priori SG, Aliot E, Blomstrom-Lundqvist C, et al. Task Force on Sudden Cardiac Death of the European Society of Cardiology. *European Heart Journal* 2001;22:1374–1450.
2. Connolly SJ. Evidence-Based Analysis of Amiodarone Efficacy and Safety. *Circulation* 1999;100:2025–2034.
3. Ezekowitz JA, Armstrong PW, McAlister A. Implantable cardioverter defibrillators in primary and secondary prevention: A systematic review of randomized, controlled trials. *Ann Intern Med* 2003;138:445–452.
4. Gregoratos G, Abrams J, Epstein AE, et al. American College of Cardiology/ American Heart Association Task Force on Practice Guidelines American College of

- Cardiology/American Heart Association/North American Society for Pacing and Electrophysiology Committee. ACC/AHA/NASPE 2002 guideline update for implantation of cardiac pacemakers and antiarrhythmia devices: Summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/NASPE Committee to Update the 1998 Pacemaker Guidelines). *Circulation* 2002;106:2145–2165.
5. Stevenson WG, Brugada P, Waldecker B, Zehender M, Wellens HJ. Clinical, angiographic, and electrophysiologic findings in patients with aborted sudden cardiac death as compared with patients with sustained ventricular tachycardia after myocardial infarction. *Circulation* 1985;71:1146–1152.
  6. Khairy P, Thibault B, Talajic M, et al. Prognostic significance of ventricular arrhythmia post-myocardial infarction. *Can J Cardiol* 2003;19:1393–1404.
  7. Newby KH, Thompson T, Stebbins A, Topol EJ, Califf RM, Natale A. Sustained ventricular arrhythmias in patients receiving thrombolytic therapy incidence and outcomes. *Circulation* 1998;98:2567–2573.
  8. Bigger JT Jr, Fleiss JL, Rolnitzky LM. Prevalence, characteristics and significance of ventricular tachycardia detected by 24-hour continuous electrocardiographic recordings in the late hospital phase of acute myocardial infarction. *Am J Cardiol* 1986;58:1151–1160.
  9. Bodegas A, Arana J, Rumoroso JR, Rodrigo D, Barrenetxea JI. Prognosis of patients with a first episode of sustained monomorphic ventricular tachycardia. *Int J Cardiol* 1998;65:181–185.
  10. Willems AR, Tijssen JG, van Capelle FJ, et al. Determinants of prognosis in symptomatic ventricular tachycardia or ventricular fibrillation late after myocardial infarction. The Dutch Ventricular Tachycardia Study Group of the Interuniversity Cardiology Institute of The Netherlands. *J Am Coll Cardiol* 1990;16:521–530.
  11. Kleiman RB, Miller JM, Buxton AE, Josephson ME, Marchlinski FE. Prognosis following sustained ventricular tachycardia occurring early after myocardial infarction. *Am J Cardiol* 1988;62:528–533.
  12. Al-Khatib SM, Stebbins AL, Califf RM, et al. Sustained ventricular arrhythmias and mortality among patients with acute myocardial infarction: Results from the GUSTO-III trial. *Am Heart J* 2003;145:515–521.
  13. Al-Khatib SM, Granger CB, Huang Y, et al. Sustained ventricular arrhythmias among patients with acute coronary syndromes with no ST-segment elevation incidence, predictors, and outcomes. *Circulation* 2002;106:309–312.
  14. Freedberg NA, Hill JN, Fogel RI, Prystowsky EN. Recurrence of symptomatic ventricular arrhythmias in patients with implantable cardioverter defibrillator after the first device therapy. *J Am Coll Cardiol* 2001;37:1910–1915.
  15. Raitt MH, Renfro EG, Epstein AE, et al. “Stable” ventricular tachycardia is not a benign rhythm: Insights from the Antiarrhythmics Versus Implantable Defibrillators (AVID) registry. *Circulation* 2001;103:244–252.
  16. Schläpfer J, Rapp F, Kappenberger L, Fromer M. Electrophysiologically guided amiodarone therapy versus the implantable cardioverter-defibrillator for sustained ventricular tachyarrhythmias after myocardial infarction results of long-term follow-up. *J Am Coll Cardiol* 2002;39:1813–1819.
  17. Sarter BH, Finkle JK, Gerszten RE, Buxton AE. What is the risk of sudden cardiac death in patients presenting with hemodynamically stable sustained ventricular tachycardia after myocardial infarction? *J Am Coll Cardiol* 1996;28:122–129.
  18. Olson PJ, Woelfel A, Simpson RJ, Foster JR. Stratification of sudden death risk in patients receiving long-term amiodarone treatment for sustained ventricular tachycardia or ventricular fibrillation. *Am J Cardiol* 1993;71:823–826.
  19. Stevenson WG, Friedman PL, Kocovic D, Sager PT, Saxon LA, Pavri B. Radiofrequency catheter ablation of ventricular tachycardia after myocardial infarction. *Circulation* 1998;98:308–314.
  20. Bocker D, Block M, Isbruch F, et al. Benefits of treatment with implantable cardioverter-defibrillators in patients with stable ventricular tachycardia without cardiac arrest. *Br Heart J* 1995;73:158–163.
  21. Raitt MH, Dolack GL, Kudenchuk PJ, Poole JE, Bardy GH. Ventricular arrhythmias detected after transvenous defibrillator implantation in patients with a clinical history of only ventricular fibrillation implications for use of implantable defibrillator. *Circulation* 1995;91:1996–2001.
  22. Caruso AC, Marcus FI, Hahn EA, Hartz VL, Mason JW. Predictors of arrhythmic death and cardiac arrest in the ES-VEM trial. Electrophysiologic Study Versus Electromagnetic Monitoring. *Circulation* 1997;96:1888–1892.
  23. The Multicenter Postinfarction Research Group. Risk stratification and survival after myocardial infarction. *N Engl J Med* 1983;309:331–336.
  24. Bigger JT Jr, Fleiss JL, Kleiger R, Miller JP, Rolnitzky LM. The relationships among ventricular arrhythmias, left ventricular dysfunction and mortality in the 2 years after myocardial infarction. *Circulation* 1984;69:250–258.
  25. Mukharji J, Rude RE, Poole WK, et al. and the MILIS Study Group. Risk factors for sudden death after acute myocardial infarction: Two-year follow-up. *Am J Cardiol* 1984;54:31–36.
  26. Coromilas J, Bigger JT Jr, Kleiger RE, and the MDPIT Research Group. Relations among left ventricular dysfunction, ventricular arrhythmias, and mortality after myocardial infarction. *Circulation* 1989;80(Suppl. II):48.
  27. Yap YG, Duong T, Bland M, et al. Left ventricular ejection fraction in the thrombolytic era remains a powerful predictor of long term but not short-term all cause cardiac and arrhythmic mortality after myocardial infarction—A secondary meta-analysis of 2828 patients. *J Am Coll Cardiol* 2001;37(Suppl. A):445A.
  28. Domanski MJ, Sakseena S, Epstein AE, et al. Relative effectiveness of the implantable cardioverter-defibrillator and antiarrhythmic drugs in patients with varying degrees of left ventricular dysfunction who have survived malignant ventricular arrhythmias. *J Am Coll Cardiol* 1999;34:1090–1095.
  29. Domanski MJ, Epstein A, Hallstrom A, Saksena S, Zipes DP. Survival of antiarrhythmic or implantable cardioverter defibrillator treated patients with varying degrees of left ventricular dysfunction who survived malignant ventricular arrhythmias. *J Cardiovasc Electrophysiol* 2002;13:580–583.
  30. CASCADE Investigators. Cardiac Arrest in Seattle: Conventional versus amiodarone drug evaluation. *Am J Cardiol* 1991;67:578–584.
  31. Weinberg BA, Miles WM, Klein LS, et al. Five-year follow-up of 589 patients treated with amiodarone. *Am Heart J* 1993;125:109–120.
  32. Ceremuzynski L, Kleczar E, Krzeminska-Pakula M, et al. Effect of amiodarone on mortality after myocardial infarction:

- A double-blind, placebo-controlled, pilot study. *J Am Coll Cardiol* 1992;20:1056-1062.
33. Cairns JA, Connolly SJ, Gent M, Roberts R. Post-myocardial infarction mortality in patients with ventricular premature depolarizations. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Pilot Study. *Circulation* 1991;84:550-557.
  34. Hockings BE, George T, Mahrous F, Taylor RR, Hajar HA. Effectiveness of amiodarone on ventricular arrhythmias during and after acute myocardial infarction. *Am J Cardiol* 1987;60:967-970.
  35. Cairns JA, Connolly SJ, Roberts R, Gent M. Randomised trial of outcome after myocardial infarction in patients with frequent or repetitive ventricular premature depolarizations: CAMIAT. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Investigators. *Lancet* 1997;349:675-682.
  36. Julian DG, Camm AJ, Frangin G, et al. Randomised trial of effect of amiodarone on mortality in patients with left-ventricular dysfunction after recent myocardial infarction: EMIAT. European Myocardial Infarct Amiodarone Trial Investigators. *Lancet* 1997;349:667-674.
  37. Sim I, McDonald KM, Lavori PW, Norbutas CM, Hlatky MA. Quantitative overview of randomized trials of amiodarone to prevent sudden cardiac death. *Circulation* 1997;96:2823-2829.
  38. Amiodarone Trials Meta-Analysis Investigators. Effect of prophylactic amiodarone on mortality after acute myocardial infarction and in congestive heart failure: Meta-analysis of individual data from 6500 patients in randomized trials. *Lancet* 1997;350:1417-1424.
  39. Bardy GH, Lee KL, Mark DB, et al. for the Sudden Death in Heart Failure Trial (SCD-HeFT) Investigators. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med* 2005;352:225-237.
  40. Waller TJ, Kay HR, Spielman SR, Kutalek SP, Greenspan AM, Horowitz LN. Reduction in sudden death and total mortality by antiarrhythmic therapy evaluated by electrophysiologic drug testing: Criteria of efficacy in patients with sustained ventricular tachyarrhythmia. *J Am Coll Cardiol* 1987;10:83-89.
  41. Horowitz LN, Greenspan AM, Spielman SR, Frye SJ, Yacone LA, Lowenthal DT. Usefulness of electrophysiologic testing in evaluation of amiodarone therapy for sustained ventricular tachyarrhythmias associated with coronary heart disease. *Am J Cardiol* 1985;55:367-371.
  42. Naccarelli GV, Fineberg NS, Zipes DP, Heger JJ, Duncan G, Prystowsky EN. Amiodarone: Risk factors for recurrence of symptomatic ventricular tachycardia identified at electrophysiologic study. *J Am Coll Cardiol* 1985;6:814-821.
  43. Zhu J, Haines DE, Lerman BB, DiMarco JP. Predictors of efficacy of amiodarone and characteristics of recurrence of arrhythmia in patients with sustained ventricular tachycardia and coronary artery disease. *Circulation* 1987;76:802-809.
  44. Kadish AH, Buxton AE, Waxman HL, Flores B, Josephson ME, Marchlinski FE. Usefulness of electrophysiologic study to determine the clinical tolerance of arrhythmia recurrences during amiodarone therapy. *J Am Coll Cardiol* 1987;10:90-96.
  45. Yazaki Y, Haffajee CI, Gold RL, Bishop RL, Alpert JS. Electrophysiologic predictors of long-term clinical outcome with amiodarone for refractory ventricular tachycardia secondary to coronary artery disease. *Am J Cardiol* 1987;60:293-297.
  46. Greenspon AJ, Volosin KJ, Greenberg RM, Jefferies L, Rotmensch HH. Amiodarone therapy: Role of early and late electrophysiologic studies. *J Am Coll Cardiol* 1988;11:117-123.
  47. Mitchell LB. The role of pharmacological therapy for ventricular tachyarrhythmias: Where to go from here? In: Singer I, Barold SS, Camm AJ, eds. *Nonpharmacological Therapy of Arrhythmias for the 21st Century: The State of the Art*. Armonk, NY: Futura Publishing Co., 1998:565-587.
  48. Credner SC, Klingenhoben T, Mauss O, Sticherling C, Hohnloser SH. Electrical storm in patients with transvenous implantable cardioverterdefibrillators: Incidence, management and prognostic implications. *J Am Coll Cardiol* 1998;32:1909-1915.
  49. Greene M, Newman D, Geist M, Paquette M, Heng D, Dorian P. Is electrical storm in ICD patients the sign of a dying heart? Outcome of patients with clusters of ventricular arrhythmias. *Europace* 2000;2:263-269.
  50. Dorian P, Mangat I. Role of Amiodarone in the Era of the Implantable Cardioverter Defibrillator. *J Cardiovasc Electrophysiol* 2003;14:S78-S81.
  51. Sra J, Bhatia A, Dhala A, et al. Electroanatomically guided catheter ablation of ventricular tachycardias causing multiple defibrillator shocks. *Pacing Clin Electrophysiol* 2001;24:1645-1652.
  52. Stevenson WG, Friedman PL, Sweeney MO. Catheter ablation as an adjunct to ICD therapy. *Circulation* 1997;96:1378-1380.
  53. Strickberger SA, Man KC, Daoud EG, et al. A prospective evaluation of catheter ablation of ventricular tachycardia as adjuvant therapy in patients with coronary artery disease and an implantable cardioverter-defibrillator. *Circulation* 1997;96:1525-1531.
  54. Willems S, Borggrefe M, Shenasa M, et al. Radiofrequency catheter ablation of ventricular tachycardia following implantation of an automatic cardioverter defibrillator. *Pacing Clin Electrophysiol* 1993;16:1684-1692.
  55. Della Bella P, De Ponti R, Uriarte JA, et al. Catheter ablation and antiarrhythmic drugs for haemodynamically tolerated post-infarction ventricular tachycardia; long-term outcome in relation to acute electrophysiological findings. *Eur Heart J* 2002;23:414-424.
  56. Connolly SJ, Hallstrom AP, Cappato R, et al. Meta-analysis of the implantable cardioverter defibrillator secondary prevention trial. *Eur Heart J* 2000;21:2071-2078.
  57. Bokhari F, Newman D, Greene M, et al. Long-term comparison of the implantable cardioverter defibrillator versus amiodarone eleven-year follow-up of a subset of patients in the Canadian implantable defibrillator study (CIDS). *Circulation* 2004;110:112-116.
  58. Salukhe TV, Dimopoulos K, Sutton R, Coats AJ, Piepoli M, Francis DP. Life-years gained from defibrillator implantation markedly nonlinear increase during 3 years of follow-up and its implications. *Circulation* 2004;109:1848-1853.
  59. Buxton AE. The clinical use of implantable cardioverter defibrillators: Where are we now? Where should we go? *Ann Intern Med* 2003;138:512-514.
  60. Davis HT, DeCamilla J, Bayer LW, Moss AJ. Survivorship patterns in the posthospital phase of myocardial infarction. *Circulation* 1979;60:1252-1258.
  61. Bigger JT Jr, Fleiss JL, Kleiger R, Miller JP, Rolnitzky LM. For the Multicenter Post-Infarction Research Group.

- The relationships among ventricular arrhythmias, left ventricular dysfunction, and mortality in the 2 years after myocardial infarction. *Circulation* 1984;69:250–258.
62. Myerburg RJ, Kessler KM, Castellanos A. Sudden cardiac death: Structure, function, and time-dependence of risk. *Circulation* 1992;85(suppl I):I-2–I-10.
  63. Wilber DJ, Zareba W, Hall J, et al. Time Dependence of Mortality Risk and Defibrillator Benefit After Myocardial Infarction. *Circulation* 2004;109:1082–1084.
  64. Exner DV, Klein GJ. Do we need a randomized trial of defibrillator therapy in every subset of patients with increased risk of sudden death? *J Cardiovasc Electrophysiol* 2003;14:574–577.
  65. Buxton AE, Lee KL, Hafley GE, et al. Relation of ejection fraction and inducible ventricular tachycardia to mode of death in patients with coronary artery disease an analysis of patients enrolled in the Multicenter Unsustained Tachycardia Trial. *Circulation* 2002;106:2466–2472.
  66. Burns RJ, Gibbons RJ, Yi Q, et al. The relationship of left ventricular ejection fraction, end-systolic volume index and infarct size to six month mortality after hospital discharge following myocardial infarction treated by thrombolysis. *J Am Coll Cardiol* 2002;39:30–36.
  67. Hallstrom A, Pratt CM, Greene HL, et al. Relations between heart failure, ejection fraction, arrhythmia suppression and mortality: Analysis of the Cardiac Arrhythmia Suppression Trial. *J Am Coll Cardiol* 1995;25:1250–1257.
  68. PCD Investigator Group. Clinical outcome of patients with malignant ventricular tachyarrhythmias and a multiprogrammable implantable cardioverter-defibrillator implanted with or without thoracotomy: An international multicenter study. *J Am Coll Cardiol* 1994;23:1521–1530.